## UNIVERSITY OF SASKATCHEWAN

## **Department of Physics and Engineering Physics**

# Physics 111.6 MIDTERM TEST #3

January 2	3, 2001		Time. 90 min	lutes
NAME:			STUDENT NO.:	
	(Last)	Please Print	(Given)	
LECTUR	E SECTION	(please circle):		
		01	Dr. R.E. Pywell	
		02	B. Zulkoskey	
		03	Dr. G.R. Davis	
		97	M. Gibson	
		C41	A. Duffy	
		C15	A Duffy	

#### **INSTRUCTIONS:**

Innuary 25, 2001

- 1. You should have a test paper, a formula sheet, and an OMR sheet. The test paper consists of 9 pages. It is the responsibility of the student to check that the test paper is complete.
- 2. Enter your name and <u>STUDENT NUMBER</u> on the OMR sheet.
- 3. The test paper, the formula sheet and the OMR sheet must all be submitted.
- 4. The test paper will be returned. The formula sheet and the OMR sheet will <u>NOT</u> be returned.

#### PLEASE DO NOT WRITE ANYTHING ON THIS TABLE

QUESTION NUMBER	MAXIMUM MARKS	MARKS OBTAINED
Part A	10	
Part B	10	
C1	5	
C2	5	
C3	5	
TOTAL	35	

Time: 00 minutes

Physics	111.6 Midterm Test #2
January	25, 2001; Page 2

### Student No.:

### PART A


FOR EACH OF THE FOLLOWING QUESTIONS IN PART A, ENTER THE MOST APPROPRIATE RESPONSE ON THE OMR SHEET.

- **A1.** Which one of the following statements is true concerning an object executing simple harmonic motion?
  - (A) Its velocity is never zero.
  - (B) Its acceleration is never zero.
  - (C) Its velocity and acceleration are zero simultaneously.
  - (D) Its velocity is zero when its acceleration is a maximum.
  - (E) Its maximum acceleration is equal to its maximum velocity.
- **A2.** The end of a car radio antenna is oscillating back and forth in simple harmonic motion. The amplitude of the oscillation is *A* and the frequency is *f*. The magnitude of the maximum acceleration is given by
  - (A)  $2\phi f^2 A$
  - (B)  $4\phi^2 f^2 A$
  - (C)  $4\phi^2 f^2 A^2$
  - (D)  $4\phi f A^2$
  - (E)  $2\phi fA$ .
- **A3.** Which one of the following statements is true for a completely enclosed fluid?
  - (A) Any change in pressure applied to the fluid produces a change in pressure elsewhere in the fluid which depends on direction.
  - (B) The pressure at all points in a fluid is independent of any pressure applied to it.
  - (C) Any change in applied pressure produces an equal change in pressure at all points within the fluid.
  - (D) An increase in pressure in one part of the fluid results in an equal decrease in pressure in another part of the fluid.
  - (E) The pressure is the same at all points within the fluid.
- **A4.** Bernoulli's Principle is a statement of
  - (A) energy conservation in dynamic fluids.
  - (B) momentum conservation in dynamic fluids.
  - (C) hydrostatic equilibrium.
  - (D) thermal equilibrium in fluids.
  - (E) mechanical equilibrium in fluids.
- **A5.** A pressure difference is maintained between two ends of a pipe of length L, and radius R, so that a viscous fluid flows from one end to the other with a flow rate Q. If the length of the pipe is increased by a factor of 16, what must the new radius of the pipe be to obtain the same flow rate Q?
  - (A) R/4
  - (B) R/2
  - (C) 2R
  - (D) 4R
  - (E) 8R

Physics	111	.6 Midterm	Test #2	2
January	25.	2001; Page	3	

Student No.:	
--------------	--

- **A6.** The tension in a taut rope is increased by a factor of 9. How does the speed of wave pulses on the rope change, if at all?
  - (A) The speed remains the same.
  - (B) The speed is reduced by a factor of 3.
  - (C) The speed is reduced by a factor of 9.
  - (D) The speed is increased by a factor of 3.
  - (E) The speed is increased by a factor of 9.
- **A7.** Which of the following superpositions will result in beats?
  - (A) The superposition of waves that travel with different speeds.
  - (B) The superposition of identical waves that travel in the same direction.
  - (C) The superposition of identical waves that travel in the opposite direction.
  - (D) The superposition of waves that are identical except for slightly different amplitudes.
  - (E) The superposition of waves that are identical except for slightly different frequencies.
- A8. Audio sound waves of the same wavelength  $\varsigma$  are emitted from two speakers that are vibrating in phase. The speakers are producing sound energy at the same rate and emitting sound uniformly in all directions. Listener A is sitting at a distance  $\varsigma$  from each of the speakers. Listener B is sitting at a distance of  $3\varsigma/2$  from one of the speakers and a distance  $\varsigma$  from the other speaker. Which one of the following statements is correct?
  - (A) Both listeners hear sound of the same loudness.
  - (B) Listener B hears sound that is slightly louder than that heard by listener A.
  - (C) Listener B hears sound that is much louder than that heard by listener A.
  - (D) Listener A hears sound that is much louder than that heard by listener B.
  - (E) The relationship between the loudness of sound heard by the listeners depends on the speed of sound.
- **A9.** Two conducting spheres, A and B, of equal size are placed on an insulating table. Sphere A has a charge of +Q and sphere B has a charge of -2Q. The spheres are allowed to touch each other and are then separated on the insulating table. Which one of the following statements is correct?
  - (A) The spheres now each carry a charge of –Q and repel each other.
  - (B) The spheres now each carry a charge of +3Q/2 and attract each other.
  - (C) The spheres now each carry a charge of -Q/2 and attract each other.
  - (D) The spheres now each carry a charge of -Q/2 and repel each other.
  - (E) Sphere A is now uncharged and sphere B has a charge of –Q.
- **A10.** Two uncharged conducting spheres, A and B, are suspended from insulating threads so that they touch each other. While a negatively-charged rod is held near, but not touching, sphere A, the two spheres are separated. How will the spheres be charged, if at all?
  - (A) Sphere A will be uncharged and sphere B will be positively charged.
  - (B) Sphere A will be negatively charged and sphere B will be positively charged.
  - (C) Both spheres will be uncharged.
  - (D) Sphere A will be negatively charged and sphere B will be uncharged.
  - (E) Sphere A will be positively charged and sphere B will be negatively charged.



•	.6 Midterm Test #2 2001; Page 4	Student No.:
PART B FOR EACH O	OF THE FOLLOWING PROBLEMS, B1 TO B5, ON PAGE PROVIDED AND ENTER YOUR ANSWERS ON PAGE 6.	ES 4 TO 6, WORK OUT THE SOLUTION IN
ONLY THE	ANSWERS WILL BE MARKED. THE SOLUTIONS WILL	NOT BE MARKED.
<b>B1.</b> How l	ong must a pendulum be if it is to have a period of	10.0 s?
0.230	is riding a moped at a constant speed of 14.0 m/s. T m and each has a moment of inertia of 0.180 kg.m <sup>2</sup> f the wheels?	The wheels of the moped have a radius of . What is the rotational kinetic energy of

continued on page 5 ...

Physics	111.6 Midterm Test #2
January	25, 2001; Page 5

**B3.** Water flows with a volume flow rate of  $1.00 \text{ m}^3/\text{s}$  in a pipe. Calculate the water speed where the pipe radius is 0.125 m.

**B4.** An ancient statue is being salvaged from the bottom of the ocean by being lifted with a cable. The mass of the statue is 70.0 kg and its volume is  $3.00 \,\Delta\,10^{42}\,\mathrm{m}^3$ . Half way to the surface the lifting is stopped temporarily. Calculate the tension in the cable as the statue, still fully submerged, is suspended motionless from the cable. The density of sea water is  $1024 \,\mathrm{kg/m}^3$ .


	ics 111.6 Midterm Test #2 ary 25, 2001; Page 6	Student No.:
B5.	A guitar string of length 0.628 $5.28 \Delta 10^{43}$ kg/m. Calculate th frequency of 165 Hz.	m, fixed at both ends, has a linear mass density $(m/L)$ of e tension that must be applied to the string to obtain a fundamental
	nequency of 100 IIE.	
ANS	WERS FOR PART B	
		T B PROBLEMS IN THE BOXES BELOW.
		EE SIGNIFICANT FIGURES AND THE UNITS MUST BE GIVEN.  KED. THE SOLUTIONS WILL NOT BE MARKED.
B1	THE MISTERS WEED DE NAME	LED. THE GOLD HONG WILL NOT BE MINNED.
B2		
В3		
<b>D</b> 4		
B4		
B5		continued on page 7
		1.48.

Janua	ary 2	11.6 Midterm Test #2 5, 2001; Page 7	Student No.:	
	ACH (	OF THE PART C QUESTIONS ON THE FOLLOWI IE FINAL ANSWER IN THE BOX PROVIDED.	NG PAGES, GIVE THE (	COMPLETE SOLUTION AND
		WERS MUST CONTAIN THREE SIGNIFICANT FIG	GURES AND THE UNITS	MUST BE GIVEN.
		<u>our Work</u> – No credit will be given for Mulae sheet must be derived.	ANSWERS ONLY. EQU	ATIONS NOT PROVIDED ON
USE	THE I	BACK OF THE PREVIOUS PAGE FOR YOUR ROU	GH WORK.	
C1.	spri disp bacl	0.500  kg air-track glider is attached to the ending constant of 20.0 N/m. Let $x = 0$ be the equal of	ilibrium position of that ilibrium position and re friction).	e glider. The glider is
	(a)	Calculate the position of the glider at time $t$	$\mid \frac{I}{8}$ .	
	(b)	Calculate the acceleration of the glider at the	e same time $t \mid \frac{T}{8}$ .	

•		11.6 Midterm Test #2 5, 2001; Page 8		Student No.:	
C2.	sou		e cyclist hears the sound with	ring away from each other. The car each frequency $f\mathcal{R}$	mits a
	(a)	Derive an expression w	which gives the speed $v_c$ in te	erms of $v, f$ and $f\mathcal{R}$	
	(b)	and the sound heard by		nitted by the car has a frequency of 4 of 387 Hz. Using your result from particular,	

continued on page 9 ...

C3. Four charges, each of magnitude  $6.00 \, \sigma C$  (two positive and two negative), are arranged at the corners of a square as shown. The sides of the square have length  $1.00 \, \text{m}$ . Calculate the electric force (magnitude and direction) acting on the charge  $q_1$  in the upper left corner. Specify the direction as the counterclockwise angle from the +x axis direction.



Magnitude:

Direction: